

Date Shipment Needed: \_\_\_\_\_ Ship To: ☐ Patient ☐ Prescriber  
☐ Nursing needed; ☐ Training needed ► All the supplies including syringes and needles will be dispensed if needed.

## RHEUMATOLOGY NON-IV REFERRAL FORM A-OL

### PATIENT INFORMATION

Patient Name:	DOB:	Sex: <input type="checkbox"/> M <input type="checkbox"/> F <input type="checkbox"/> Other:	Weight:	<input type="checkbox"/> lbs. <input type="checkbox"/> kg.
SSN:	Phone:	Allergies:		
Address:	City:	State:	Zip:	
Emergency Contact:	Phone:	<input type="checkbox"/> Additional Information Attached		

### PRESCRIBER INFORMATION

Prescriber:	NPI:	DEA:	State Lic:
Supervising Physician:	Practice Name:		
Address:	City:	State:	Zip:
Phone:	Fax:	Key Office Contact:	Phone:

### DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT

**Primary Diagnosis:** ☐ M06.9 Rheumatoid Arthritis ☐ L40.54; L40.59 Psoriatic Arthritis ☐ M08.00 Unspecified Juvenile Rheumatoid Arthritis ☐ M08.3 Juvenile Rheumatoid Polyarthrits (Seronegative)  
☐ M08.20 Juvenile Idiopathic Arthritis ☐ M45.9 Ankylosing Spondylitis ☐ M33.20 Polymyositis ☐ M81.0 Osteoporosis ☐ M15.0; M15.9 Osteoarthritis ☐ Other: \_\_\_\_\_

• Has patient been treated *previously* for this condition? ☐ Yes ☐ No Is patient *currently* on therapy? ☐ Yes ☐ No Please list medication(s) and treatment duration: \_\_\_\_\_

• Will patient stop taking the above medication(s) before starting the new medication? ☐ Yes ☐ No If yes, how long should patient wait before starting the new medication? \_\_\_\_\_

• Other medications patient is currently taking including OTC medications with dosage and direction (or fax medication profile): \_\_\_\_\_

• Has patient received a **Quatiferon gold**, **Tspot** or **PPD (tuberculosis) Skin Test**? ☐ Yes ☐ No Date: \_\_\_\_\_ Results: ☐ Negative ☐ Positive  
*Prior to initiating treatment and periodically during therapy, patient should be evaluated for active tuberculosis and tested for latent infection*

### INSURANCE INFORMATION

☐ Please attach front and back of patient's insurance card (medical and prescription)

### COPAY CARD ENROLLMENT

☐ Please check if enrolling in copay card Copay ID: \_\_\_\_\_

### PRESCRIPTION INFORMATION

<input type="checkbox"/> <b>Actemra® 162 mg</b> <input type="checkbox"/> Pen <input type="checkbox"/> Prefilled Syringe <input type="checkbox"/> <100 kg: 162 mg SQ once every other week <input type="checkbox"/> ≥100 kg: 162 mg SQ once every week <input type="checkbox"/> <b>Cimzia® 200 mg/mL Prefilled Syringe</b> <input type="checkbox"/> <b>Cimzia® 200 mg Vial</b> <input type="checkbox"/> Starter Dose Not Needed <small>*Cimzia vial should be prepared and administered by a healthcare professional. Prefilled Syringe will be dispensed unless vial is requested.</small> <b>Starter Dose:</b> <input type="checkbox"/> 400 mg SQ (2 inj. of 200 mg) initially (Week 0), repeat at Weeks 2 and 4 <b>Maintenance Dose:</b> <input type="checkbox"/> 400 mg SQ (2 inj. of 200 mg) every 4 weeks <input type="checkbox"/> 200 mg SQ every 2 weeks <b>Alternate Dose:</b> <input type="checkbox"/> _____	QTY: <u>2 Pens / Syringes</u>   Refills: _____ QTY: <u>4 Pens / Syringes</u>   Refills: _____ <input type="checkbox"/> <b>Enroll in Cimplicity™ Program</b> QTY: <u>1 Starter Kit (6 PFS)</u>   Refills: <u>0</u> QTY: <u>1 Box (2 INJ.)</u>   Refills: _____ QTY: _____   Refills: _____
<input type="checkbox"/> <b>Cosentyx® 150 mg/mL Prefilled Syringe</b> <input type="checkbox"/> <b>Cosentyx® 150 mg/mL Sensoready Pen</b> <input type="checkbox"/> <b>Cosentyx® 300 mg/2 mL UnoReady Pen</b> <b>Starter Dose:</b> <input type="checkbox"/> 150 mg SQ at Weeks 0, 1, 2, 3 <input type="checkbox"/> 300 mg SQ at Weeks 0, 1, 2, 3 <input type="checkbox"/> Starter Dose Not Needed <b>Maintenance Dose:</b> <input type="checkbox"/> 150 mg SQ every 4 weeks (starting at Week 4) <input type="checkbox"/> 300 mg SQ every 4 weeks (starting at Week 4)	QTY: <u>QS 28 Day Supply</u>   Refills: <u>0</u> QTY: <u>QS 28 Day Supply</u>   Refills: _____
<input type="checkbox"/> <b>Enbrel® 50 mg/mL Sureclick (Autoinjector)</b> <input type="checkbox"/> <b>Enbrel® 50 mg/mL Prefilled Syringe</b> <small>*Not to be used in pediatric weighing less than 63 kg (138 lb.)</small> <input type="checkbox"/> 50 mg SQ weekly <input type="checkbox"/> <b>Alternate Dose:</b> _____	<input type="checkbox"/> <b>Enroll in Enliven® Program</b> QTY: <u>4</u>   Refills: _____ QTY: _____   Refills: _____
<input type="checkbox"/> <b>Enbrel® 25 mg/0.5 mL Prefilled Syringe</b> <input type="checkbox"/> 25 mg SQ twice weekly (72-96 hours apart) <input type="checkbox"/> <b>Alternate Dose:</b> _____	QTY: <u>8</u>   Refills: _____ QTY: _____   Refills: _____
<input type="checkbox"/> <b>Humira® 40 mg/0.4 mL Pen CF</b> <input type="checkbox"/> <b>Humira® 40 mg/0.4 mL Prefilled Syringe CF</b> <input type="checkbox"/> Inject 40 mg SQ every other week <input type="checkbox"/> Inject 40 mg SQ every week	<input type="checkbox"/> <b>Enroll in Humira Complete Program</b> QTY: _____   Refills: _____
<input type="checkbox"/> <b>Humira® 10 mg/0.2 mL Prefilled Syringe CF</b> <input type="checkbox"/> <b>Humira® 20 mg/0.4 mL Prefilled Syringe CF</b> <input type="checkbox"/> Inject 10 mg SQ every other week <input type="checkbox"/> Inject 20 mg SQ every other week	QTY: _____   Refills: _____ QTY: _____   Refills: _____
<input type="checkbox"/> <b>Ilaris® 150 mg SDV Inj.</b> _____ mg SQ every _____ Weeks	QTY: _____   Refills: _____
<input type="checkbox"/> <b>Ilaris® 150 mg/mL Single Dose Vial</b> <input type="checkbox"/> Inject 4 mg/kg _____ mg SQ every 4 weeks (300 mg/dose maximum)	QTY: _____   Refills: _____
<input type="checkbox"/> <b>Kevzara® Inj. Single Prefilled Syringe</b> <input type="checkbox"/> <b>Kevzara® Inj. Single Prefilled Pen</b> <input type="checkbox"/> 150 mg/1.14 mL <input type="checkbox"/> 200 mg/1.14 mL <input type="checkbox"/> 1 SQ inj. every 2 weeks	QTY: _____   Refills: _____
<input type="checkbox"/> <b>Olumiant® 2 mg Tablet</b> <input type="checkbox"/> <b>1 mg Tablet</b> <input type="checkbox"/> Take 2 mg tablet PO once daily <input type="checkbox"/> Take 1 mg tablet PO once daily	QTY: _____   Refills: _____ QTY: _____   Refills: _____

**Prescriber's Signature:** \_\_\_\_\_ ☐ DAW (Dispense as Written) **Date:** \_\_\_\_\_

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. NO STAMPED SIGNATURES WILL BE ACCEPTED. Where required by law, send electronic prescription or on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

**IMPORTANT NOTICE:** This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.

Date Shipment Needed: \_\_\_\_\_ Ship To: ☐ Patient ☐ Prescriber  
☐ Nursing needed; ☐ Training needed ► All the supplies including syringes and needles will be dispensed if needed.

## RHEUMATOLOGY NON-IV REFERRAL FORM OM-Z

### PATIENT INFORMATION

Patient Name:	DOB:	Sex: <input type="checkbox"/> M <input type="checkbox"/> F <input type="checkbox"/> Other:	Weight:	<input type="checkbox"/> lbs. <input type="checkbox"/> kg.
SSN:	Phone:	Allergies:		
Address:	City:	State:	Zip:	
Emergency Contact:	Phone:	<input type="checkbox"/> Additional Information Attached		

### PRESCRIBER INFORMATION

Prescriber:	NPI:	DEA:	State Lic:
Supervising Physician:	Practice Name:		
Address:	City:	State:	Zip:
Phone:	Fax:	Key Office Contact:	Phone:

### DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT

**Primary Diagnosis:** ☐ M06.9 Rheumatoid Arthritis ☐ L40.54; L40.59 Psoriatic Arthritis ☐ M08.00 Unspecified Juvenile Rheumatoid Arthritis ☐ M08.3 Juvenile Rheumatoid Polyarthritits (Seronegative)  
☐ M08.20 Juvenile Idiopathic Arthritis ☐ M45.9 Ankylosing Spondylitis ☐ M33.20 Polymyositis ☐ M81.0 Osteoporosis ☐ M15.0; M15.9 Osteoarthritis ☐ Other: \_\_\_\_\_

• Has patient been treated *previously* for this condition? ☐ Yes ☐ No Is patient *currently* on therapy? ☐ Yes ☐ No Please list medication(s) and treatment duration: \_\_\_\_\_

• Will patient stop taking the above medication(s) before starting the new medication? ☐ Yes ☐ No If yes, how long should patient wait before starting the new medication? \_\_\_\_\_

• Other medications patient is currently taking including OTC medications with dosage and direction (or fax medication profile): \_\_\_\_\_

• Has patient received a **Quatiferon gold**, **Tspot** or **PPD (tuberculosis) Skin Test**? ☐ Yes ☐ No Date: \_\_\_\_\_ Results: ☐ Negative ☐ Positive  
*Prior to initiating treatment and periodically during therapy, patient should be evaluated for active tuberculosis and tested for latent infection*

### INSURANCE INFORMATION

☐ Please attach front and back of patient's insurance card (medical and prescription)

### COPAY CARD ENROLLMENT

☐ Please check if enrolling in copay card **Copay ID:** \_\_\_\_\_

### PRESCRIPTION INFORMATION

<input type="checkbox"/> <b>Otezla® Tablet</b> <input type="checkbox"/> Five (5) day titration period: <b>Day 1:</b> 10 mg, <b>Day 2:</b> 10 mg BID, <b>Day 3:</b> 10 mg in AM then 20 mg in PM, <b>Day 4:</b> 20 mg BID, <b>Day 5:</b> 20 mg in AM then 30 mg in PM <input type="checkbox"/> After five (5) day titration period, 30 mg BID	QTY: _____ 1 Kit   Refills: _____ 0
<input type="checkbox"/> <b>Orencia® 125 mg Prefilled Syringe</b> <input type="checkbox"/> <b>Orencia® 125 mg ClickJect Autoinjector</b> <input type="checkbox"/> Starter Dose: One dose of IV infusion (per body weight) <input type="checkbox"/> <60 kg: 500 mg IV x 1 dose <input type="checkbox"/> 60 - 100 kg: 750 mg IV x 1 dose <input type="checkbox"/> > 100 kg: 1000 mg IV x 1 dose <input type="checkbox"/> Maintenance Dose: 150 mg SQ every week	QTY: _____ 60 Tablets   Refills: _____ <input type="checkbox"/> <b>Enroll in Orencia OnCall Program</b> QTY: _____ 2 x 250 mg Vial   Refills: _____ 0 QTY: _____ 3 x 250 mg Vial   Refills: _____ 0 QTY: _____ 4 x 250 mg Vial   Refills: _____ 0 QTY: _____ 4 PFS / Pens   Refills: _____
<input type="checkbox"/> <b>Rinvoq® 15 mg Oral Tablet</b> Take one tablet orally once daily with or without food	QTY: _____ 30   Refills: _____
<input type="checkbox"/> <b>Siliq® 210 mg / 1.5 mL Prefilled Syringe</b> <input type="checkbox"/> 210 mg SQ at Weeks 0, 1, 2 <input type="checkbox"/> 210 mg SQ every 2 weeks (starting at week 2)	QTY: _____ 2 Pens   Refills: _____ 0 QTY: _____ 2 Pens   Refills: _____
<input type="checkbox"/> <b>Skyrizi® 150 mg/mL Pen</b> <input type="checkbox"/> <b>Skyrizi® 150 mg/mL Prefilled Syringe</b> <input type="checkbox"/> Inj. 150 mg SQ at Week 0 <input type="checkbox"/> Inj. 150 mg SQ every 12 weeks (starting at Week 4)	QTY: _____ 1   Refills: _____ 0 QTY: _____ 1   Refills: _____
<input type="checkbox"/> <b>Simponi® 50 mg / 0.5 mL SmartJect (Autoinjector)</b> <input type="checkbox"/> <b>Simponi® 50 mg / 0.5 mL Prefilled Syringe</b> <input type="checkbox"/> 50 mg SQ every month <input type="checkbox"/> Alternate Dose: _____	<input type="checkbox"/> <b>Enroll in SimponiOne Program</b> QTY: _____   Refills: _____ QTY: _____   Refills: _____
<input type="checkbox"/> <b>Taltz® 80 mg/mL</b> <input type="checkbox"/> <b>Pen</b> <input type="checkbox"/> <b>Prefilled Syringe</b> <input type="checkbox"/> Starter Dose Not Needed Starter Dose: <input type="checkbox"/> Inject 160 mg once SQ on Week 0 Maintenance Dose: <input type="checkbox"/> Inject 80 mg once SQ every 4 weeks	QTY: _____ 2   Refills: _____ QTY: _____ 1 Pen / 1 Syringe   Refills: _____
<input type="checkbox"/> <b>Tremfya® 100 mg/mL</b> <input type="checkbox"/> <b>Pen</b> <input type="checkbox"/> <b>Prefilled Syringe</b> <input type="checkbox"/> Starter Dose Not Needed Starter Dose: <input type="checkbox"/> 100 mg SQ at Week 0 and Week 4 Maintenance Dose: <input type="checkbox"/> Inject 80 mg SQ once every 4 weeks (starting at Week 4)	QTY: _____ 1 Pen / 1 Syringe   Refills: _____ 0 QTY: _____ 1 Pen / 1 Syringe   Refills: _____
<input type="checkbox"/> <b>Xeljanz® 5 mg Tablet</b> <input type="checkbox"/> 5 mg PO BID	QTY: _____   Refills: _____
<input type="checkbox"/> <b>Xeljanz® XR 11 mg Tablet</b> <input type="checkbox"/> 11 mg PO once daily	QTY: _____   Refills: _____
<input type="checkbox"/> <b>Other:</b> _____	QTY: _____   Refills: _____

**Prescriber's Signature:** \_\_\_\_\_ ☐ DAW (Dispense as Written)

**Date:** \_\_\_\_\_

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. NO STAMPED SIGNATURES WILL BE ACCEPTED. Where required by law, send electronic prescription or on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

**IMPORTANT NOTICE:** This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.