

Date Shipment Needed: \_\_\_\_\_ Ship To:  Patient  Prescriber  
 Nursing needed;  Training needed ► All the supplies including syringes and needles will be dispensed if needed.

**RHEUMATOLOGY IV ROUTE REFERRAL FORM**

**PATIENT INFORMATION**

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Sex:  M  F  Other: \_\_\_\_\_ Weight: \_\_\_\_\_  lbs.  kg.  
 SSN: \_\_\_\_\_ Phone: \_\_\_\_\_ Allergies: \_\_\_\_\_  
 Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_  
 Emergency Contact: \_\_\_\_\_ Phone: \_\_\_\_\_  **Additional Information Attached**

**PRESCRIBER INFORMATION**

Prescriber: \_\_\_\_\_ NPI: \_\_\_\_\_ DEA: \_\_\_\_\_ State Lic: \_\_\_\_\_  
 Supervising Physician: \_\_\_\_\_ Practice Name: \_\_\_\_\_  
 Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_  
 Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Key Office Contact: \_\_\_\_\_ Phone: \_\_\_\_\_

**DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT**

**Primary Diagnosis:**  M06.9 Rheumatoid Arthritis  L40.54; L40.59 Psoriatic Arthritis  M08.00 Unspecified Juvenile Rheumatoid Arthritis  M08.20 Juvenile Idiopathic Arthritis  
 M08.3 Juvenile Rheumatoid Polyarthritits (Seronegative)  M45.9 Ankylosing Spondylitis  M33.20 Polymyositis  M15.0 - M15.9 Osteoarthritis  Other: \_\_\_\_\_  
 • Has patient been treated *previously* for this condition?  Yes  No Is patient *currently* on therapy?  Yes  No Please list medication(s) and treatment duration: \_\_\_\_\_  
 • Will patient stop taking the above medication(s) before starting the new medication?  Yes  No If yes, how long should patient wait before starting the new medication? \_\_\_\_\_  
 • Other medications patient is currently taking including OTC medications with dosage and direction (or fax medication profile): \_\_\_\_\_  
 • Has patient received a **Quatiferon gold, Tspot or PPD (tuberculosis) Skin Test**?  Yes  No Date: \_\_\_\_\_ Results:  Negative  Positive  
*Prior to initiating treatment and periodically during therapy, patient should be evaluated for active tuberculosis and tested for latent infection*

**INSURANCE INFORMATION**

**Please attach front and back of patient's insurance card (medical and prescription)**

**COPY CARD ENROLLMENT**

**Please check if enrolling in copay card** Copay ID: \_\_\_\_\_

**PRESCRIPTION INFORMATION**

**STC Standard Protocol** will include the following: (1) dispensing ordered med/dose, (2) diluent to mix and / or dilute dose, (3) flushes to flush line and anakit med (epinephrine 0.3 mg IM / 0.15 mg IM (for pediatric patients) and diphenhydramine 50 mg/mL) and (4) premeds to take 30 mins before orally (Apap 325 mg, may repeat x1, and diphenhydramine 25 mg, may repeat x1).

**Actemra® 20 mg/mL (Vial Sizes: 4 mL, 10 mL, 20 mL) \*Maximum dose per infusion is 800 mg**  Starter Dose Not Needed  **Enroll in Access Solutions**  
 Standard Dose: 4 mg/kg \_\_\_\_\_ mg IV every 4 weeks, infusion over 60 minutes (infusion at MD's office or infusion center) QTY: \_\_\_\_\_ QS | Refills: \_\_\_\_\_  
 Increased Dose: 8 mg/kg \_\_\_\_\_ mg IV every 4 weeks, infusion over 60 minutes (infusion at MD's office or infusion center) QTY: \_\_\_\_\_ QS | Refills: \_\_\_\_\_  
 Alternate Dose: \_\_\_\_\_ QTY: \_\_\_\_\_ | Refills: \_\_\_\_\_

**IVIG**  
 Home Infusion (*Download IVIG referral form at www.AcariaHealth.com, or ask your local AcariaHealth rep*)

**Orencia® 250 mg IV Vial** \*Dosage based on patient's weight  MD's office infusion  Infusion supplies needed  **Enroll in The Circle Program**  
 Infuse \_\_\_\_\_ mg at Weeks 0, 2 and 4, then every 4 weeks thereafter QTY: \_\_\_\_\_ | Refills: \_\_\_\_\_

**Remicade® and Remicade® Biosimilars ( Inflectra®, Renflexis®, Avsola® )**  
 **Remicade® 100 mg vial**  **Inflectra® 100 mg vial**  **Renflexis® 100 mg vial**  **Avsola® 100 mg vial**  **Enroll in AccessOneSM Program**  
 MD's Office Infusion  Infusion Supplies Required QTY: QS for Approp. Amt. | Refills: 0  
 Infuse \_\_\_\_\_ mg IV on Weeks 0, 2, and 6 QTY: QS for Approp. Amt. | Refills: \_\_\_\_\_  
 Infuse \_\_\_\_\_ mg IV every 8 weeks QTY: QS for Approp. Amt. | Refills: \_\_\_\_\_  
 Infuse \_\_\_\_\_ mg IV every \_\_\_\_\_ weeks

**Rituxan® and Rituxan® Biosimilars ( Truxima®, Ruxience®, Riabni® )**  
 **Rituxan® 500 mg vial**  **Truxima® 500 mg vial**  **Ruxience® 500 mg vial**  **Riabni® 500 mg vial**  **Enroll in RISE Program**  
 Infuse 1000 mg IV once every 2 weeks for 2 doses, every 16 to 24 weeks QTY: \_\_\_\_\_ 4 Vials | Refills: \_\_\_\_\_  
 Other: \_\_\_\_\_ QTY: \_\_\_\_\_ QS | Refills: \_\_\_\_\_

**Simponi® Aria 50 mg/4 mL Single-use Vial**  Starter Dose Not Needed  
 Starter Dose: 2 mg/kg \_\_\_\_\_ mg IV on Weeks 0 and 4, infusion over 30 minutes QTY: \_\_\_\_\_ QS | Refills: \_\_\_\_\_  
 Maintenance Dose: 2 mg/kg \_\_\_\_\_ mg IV every 8 weeks, infusion over 30 minutes QTY: \_\_\_\_\_ QS | Refills: \_\_\_\_\_  
 Alternate Dose: \_\_\_\_\_ QTY: \_\_\_\_\_ QS | Refills: \_\_\_\_\_

**Other:** \_\_\_\_\_ QTY: \_\_\_\_\_ | Refills: \_\_\_\_\_

**Prescriber's Signature:** \_\_\_\_\_  DAW (Dispense as Written) **Date:** \_\_\_\_\_

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. **NO STAMPED SIGNATURES WILL BE ACCEPTED.** Where required by law, send electronic prescription or on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

**IMPORTANT NOTICE:** This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.