

Date Shipment Needed: \_\_\_\_\_ Ship To:  Patient  Prescriber

Nursing needed;  Training needed ► All the supplies including syringes and needles will be dispensed if needed.

**DERMATOLOGY REFERRAL FORM ( A - D )**

**PATIENT INFORMATION**

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Sex:  M  F  Other: \_\_\_\_\_ Weight: \_\_\_\_\_  lbs.  kg.

SSN: \_\_\_\_\_ Phone: \_\_\_\_\_ Allergies: \_\_\_\_\_

Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_

Emergency Contact: \_\_\_\_\_ Phone: \_\_\_\_\_  Additional Information Attached

**PRESCRIBER INFORMATION**

Prescriber: \_\_\_\_\_ NPI: \_\_\_\_\_ DEA: \_\_\_\_\_ State Lic: \_\_\_\_\_

Supervising Physician: \_\_\_\_\_ Practice Name: \_\_\_\_\_

Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_

Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Key Office Contact: \_\_\_\_\_ Phone: \_\_\_\_\_

**DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT**

**Primary Diagnosis:**  L20, L20.8, L20.9 Atopic Dermatitis  L28.1 Prurigo nodularis  L40.0 Psoriasis  L40.1; L40.2; L40.3, L40.4, L40.8, L40.54 Psoriatic arthritis  L40.59  L50.1 Chronic Idiopathic Urticaria  L73.2 Hidradenitis Suppurativa  Other: \_\_\_\_\_

▪ Location:  Hands  Feet  Face  Scalp  Groin  Nails  Other: \_\_\_\_\_

▪ Severity:  Mild (up to 3% BSA)  Moderate (3-10% BSA)  Severe (greater than 10% BSA), BSA \_\_\_\_\_ %

If treated previously for this condition, please indicate which drugs have been tried and failed: \_\_\_\_\_

Date range of previous therapy: \_\_\_\_\_

▪ Is patient currently on therapy?  Yes  No Type / medication(s): \_\_\_\_\_

▪ Will patient stop taking the above medication(s) before starting the new medication?  Yes  No, if yes, how long should patient wait before starting the new medication? \_\_\_\_\_

▪ Has patient received a PPD (tuberculosis) Skin Test?  Yes  No Results: \_\_\_\_\_

Prior to initiating treatment and periodically during therapy, patient should be evaluated for active tuberculosis and tested for latent infection.

**INSURANCE INFORMATION**

Please attach front and back of patient's insurance card (medical and prescription)

**COPAY CARD ENROLLMENT**

Please check if enrolling in copay card **Copay ID:** \_\_\_\_\_

**PRESCRIPTION INFORMATION**

**Adbry®**  150 mg Prefilled Syringe

Starter Dose: 600 mg SQ on Day 1  Starter Dose not needed QTY: #4 Syringes / 14 DS | Refills: 0

After 16 wks of treatment  Maintenance Dose: 300 mg SQ every 4 weeks QTY: #2 Syringes / 28 DS | Refills: \_\_\_\_\_

Maintenance Dose: 300 mg SQ every other week QTY: #4 Syringes / 28 DS | Refills: \_\_\_\_\_

**Adbry®**  300 mg Pen

Starter Dose: 600 mg SQ on Day 1  Starter Dose not needed QTY: #2 Pens / 14 DS | Refills: 0

After 16 wks of treatment  Maintenance Dose: 300 mg SQ every 4 weeks QTY: #1 Pens / 28 DS | Refills: \_\_\_\_\_

Maintenance Dose: 300 mg SQ every other week QTY: #2 Pens / 28 DS | Refills: \_\_\_\_\_

**Adbry®**  150 mg Prefilled Syringe (Pediatric: 12 years and older)

Starter Dose: 300 mg SQ on Day 1  Starter Dose not needed QTY: #2 Syringes / 14 DS | Refills: 0

Maintenance Dose: 150 mg SQ every other week QTY: #2 Syringes / 28 DS | Refills: \_\_\_\_\_

**Bimzelx®**  160 mg/mL Pen OR  160 mg/mL Syringe

320 mg (given as two 160 mg injections) SQ every 4 weeks for the first 16 weeks QTY: 2 | Refills: 4

320 mg (given as two 160 mg injections) SQ every 8 weeks QTY: 2 | Refills: \_\_\_\_\_

**Cibinqo®**  50 mg Tablet  100 mg Tablet  200 mg Tablet

1 tablet PO once daily  Other: \_\_\_\_\_ QTY: 1 Month | Refills: \_\_\_\_\_

**Cimzia®**  200 mg Vial  200 mg/mL Prefilled Syringe

400 mg/mL SQ every 2 weeks

400 mg SQ at Weeks 0, 2, 4, then 200 mg every other week thereafter (patient ≤ 90 kg) QTY: 1 Month | Refills: \_\_\_\_\_

**Cosentyx®**  150 mg/mL Sensoready® Pen  150 mg/mL Prefilled Syringe  300 mg UnoReady Pen

\*Sensoready® Pen will be dispensed if no preference indicated

Starter Dose: 300 mg SQ initially (Weeks 0, 1, 2, 3 and 4) then 300 mg SQ every 4 weeks thereafter (Week 4)  Starter Dose not needed QTY: 5 Weeks | Refills: 0

Maintenance Dose: 300 mg SQ every 4 weeks QTY: 1 Month | Refills: \_\_\_\_\_

Other: \_\_\_\_\_ QTY: 1 Month | Refills: \_\_\_\_\_

**Dupixent®**  200 mg Pen Autoinjector  200 mg Prefilled Syringe  300 mg Pen Autoinjector  300 mg Prefilled Syringe

(Dupilumab) \*Pen will be dispensed if no preference indicated for adult dosing. Prefilled Syringe may be used in ages ≥ 6 months. Prefilled Pen is only for use in ages ≥ 2 years.

**Adults:**  Starter Dose: Inj. 600 mg SQ on Day 1, then 300 mg SQ every 2 weeks starting on Day 15  Starter Dose not needed QTY: QS for Starter | Refills: 0

Maintenance Dose: Inj. 300 mg SQ every 2 Weeks QTY: 1 Month | Refills: \_\_\_\_\_

**Infants & Children:** ≥ 6 mo - < 6 yrs: Initial loading dose not necessary in pediatric patients < 6 yrs.

5 to < 15 kg: Dupixent 200 mg SQ every 4 weeks  15 to < 30 kg: Dupixent 300 mg SQ every 4 weeks QTY: 1 Box: 2 Pens/Syringes | Refills: \_\_\_\_\_

**Children & Adolescents:** ≥ 6 years - ≤ 17 years:

15 to < 30 kg: Initial: 600 mg SQ once (administered as two 300 mg injections) QTY: 1 Box: 2 Pens/Syringes | Refills: 0

15 to < 30 kg: Maintenance: 300 mg SQ every 4 weeks QTY: 1 Box: 2 Pens/Syringes | Refills: \_\_\_\_\_

30 to < 60 kg: Initial: 400 mg SQ once (administered as two 200 mg injections) QTY: 1 Box: 2 Pens/Syringes | Refills: 0

30 to < 60 kg: Maintenance: 200 mg SQ every other week QTY: 1 Box: 2 Pens/Syringes | Refills: \_\_\_\_\_

≥ 60 kg: Initial: 600 mg SQ once (administered as two 300 mg injections) QTY: 1 Box: 2 Pens/Syringes | Refills: 0

≥ 60 kg: Maintenance: 300 mg SQ every other week QTY: 1 Box: 2 Pens/Syringes | Refills: \_\_\_\_\_

**Prescriber's Signature:** \_\_\_\_\_  DAW (Dispense as Written) **Date:** \_\_\_\_\_

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. NO STAMPED SIGNATURES WILL BE ACCEPTED. Where required by law, send electronic prescription or on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

**IMPORTANT NOTICE:** This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.

Date Shipment Needed: \_\_\_\_\_ Ship To:  Patient  Prescriber

Nursing needed;  Training needed ► All the supplies including syringes and needles will be dispensed if needed.

**DERMATOLOGY REFERRAL FORM ( E - K )**

**PATIENT INFORMATION**

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Sex:  M  F  Other: \_\_\_\_\_ Weight: \_\_\_\_\_  lbs.  kg.

SSN: \_\_\_\_\_ Phone: \_\_\_\_\_ Allergies: \_\_\_\_\_

Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_

Emergency Contact: \_\_\_\_\_ Phone: \_\_\_\_\_  Additional Information Attached

**PRESCRIBER INFORMATION**

Prescriber: \_\_\_\_\_ NPI: \_\_\_\_\_ DEA: \_\_\_\_\_ State Lic: \_\_\_\_\_

Supervising Physician: \_\_\_\_\_ Practice Name: \_\_\_\_\_

Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_

Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Key Office Contact: \_\_\_\_\_ Phone: \_\_\_\_\_

**DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT**

**Primary Diagnosis:**  L20, L20.8, L20.9 Atopic Dermatitis  L28.1 Prurigo nodularis  L40.0 Psoriasis  L40.1; L40.2; L40.3, L40.4, L40.8, L40.54 Psoriatic arthritis  L40.59  
 L50.1 Chronic Idiopathic Urticaria  L73.2 Hidradenitis Suppurativa  Other: \_\_\_\_\_

• Location:  Hands  Feet  Face  Scalp  Groin  Nails  Other: \_\_\_\_\_

• Severity:  Mild (up to 3% BSA)  Moderate (3-10% BSA)  Severe (greater than 10% BSA), BSA \_\_\_\_\_ %

If treated previously for this condition, please indicate which drugs have been tried and failed: \_\_\_\_\_

Date range of previous therapy: \_\_\_\_\_

• Is patient currently on therapy?  Yes  No Type / medication(s): \_\_\_\_\_

• Will patient stop taking the above medication(s) before starting the new medication?  Yes  No, if yes, how long should patient wait before starting the new medication? \_\_\_\_\_

• Has patient received a PPD (tuberculosis) Skin Test?  Yes  No Results: \_\_\_\_\_

Prior to initiating treatment and periodically during therapy, patient should be evaluated for active tuberculosis and tested for latent infection.

**INSURANCE INFORMATION**

Please attach front and back of patient's insurance card (medical and prescription)

**COPAY CARD ENROLLMENT**

Please check if enrolling in copay card **Copay ID:** \_\_\_\_\_

**PRESCRIPTION INFORMATION**

**Ebglyss™**  250 mg / 2 mL Pen  250 mg / 2 mL Prefilled Syringe

Starter Dose: 500 mg (2 injections) SQ at Weeks 0 and 2  Starter Dose not needed QTY:      QS 28 DS | Refills:      0

Maintenance Dose: 250 mg (1 injection) SQ every 2 weeks until Week 16 or later QTY:      QS 84 DS | Refills:     

Optional: 250 mg (1 injection) SQ every 4 weeks QTY:      QS 28 DS | Refills:     

**Enbrel®**  50 mg / mL SureClick (Autoinjector)  50 mg Prefilled Syringe  Mini 50 mg Cartridge \*SureClick will be dispensed if no preference indicated  Enroll in Enbren® Program

Starter Dose: 50 mg SQ twice weekly (72 - 96 hours apart) for 3 months  Starter Dose not needed QTY:      1 Month | Refills:      2

Maintenance Dose: 50 mg SQ weekly  Other: \_\_\_\_\_ QTY:      1 Month | Refills:     

**Enbrel®**  25 mg / 0.5 mL Prefilled Syringe  25 mg Single-Use Vial \*Prefilled Syringe will be dispensed if no preference indicated

25 mg SQ twice weekly (72 - 96 hours apart)  Other: \_\_\_\_\_ QTY:      1 Month | Refills:     

**Erivedge®**  150 mg Capsules Take 1 capsule orally once daily QTY:      28 Capsules | Refills:     

**Humira®**  CF Pen Psoriasis Starter Kit NDC: 0074-1539-03  CF 40 mg / 0.4 mL Prefilled Syringe NDC: 0074-0243-02  Enroll in Humira Complete Program

\*Pen Starter Kit will be dispensed if no preference indicated

Starter Dose  One 80 mg SQ inj. Day 1, one 40 mg SQ inj. Day 8, one 40 mg SQ inj. Day 22 (OR)  Starter Dose not needed QTY:      3 Pens | Refills:      0

for Psoriasis:  Two 40 mg SQ inj. Day 1, one 40 mg SQ inj. Day 8, one 40 mg SQ inj. Day 22 QTY:      4 Syringes | Refills:      0

**Humira®**  CF 40 mg / 0.4 mL Pen NDC: 0074-0554-02  CF 40 mg / 0.4 mL Syringe NDC: 0074-0243-02

\*Pen will be dispensed if no preference indicated

Maintenance Dose for Psoriasis: 40 mg SQ once every other week QTY:      1 Month | Refills:     

**Humira®**  Starter Pkg CF 80 mg / 0.8 mL Pen NDC: 0074-0124-03  CF 40 mg / 0.4 mL Prefilled Syringe NDC: 0074-0243-02

\*Pen will be dispensed if no preference indicated

Starter Dose for  Inj. 160 mg SQ Day 1, then 80 mg SQ Day 15 (OR)  Starter Dose not needed QTY:      1 Month | Refills:      0

Hidradenitis Suppurativa:  Inj. 80 mg SQ Day 1, and 80 mg SQ Day 2, then 80 mg SQ Day 15 QTY:      1 Month | Refills:      0

**Humira®**  CF 40 mg / 0.4 mL Pen NDC: 0074-0554-02  CF 40 mg / 0.4 mL Syringe NDC: 0074-0243-02

\*Pen will be dispensed if no preference indicated

Maintenance Dose for Hidradenitis Suppurativa: 40 mg SQ Day 29 and every week thereafter QTY:      1 Month | Refills:     

Other: \_\_\_\_\_ QTY:      | Refills:     

**Ilumya®**  100 mg / mL Prefilled Syringes

Starter Dose: 100 mg SQ on Week 0 and Week 4  Starter Dose not needed QTY:      1 Month (1 PFS) | Refills:      0

Maintenance Dose: 100 mg SQ every 12 weeks (starting at Week 4) QTY:      1 Syringe | Refills:     

**Kezvara®**  200 mg / 1.14 mL Pen Autoinjector  200 mg / 1.14 mL Prefilled Syringe \*Pens will be dispensed if no preference is indicated

(Sarilumab)  200 mg subcutaneously every 2 Weeks QTY:      1 Box (2) | Refills:     

**Prescriber's Signature:** \_\_\_\_\_  DAW (Dispense as Written) **Date:** \_\_\_\_\_

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. NO STAMPED SIGNATURES WILL BE ACCEPTED. Where required by law, send electronic prescription or on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

**IMPORTANT NOTICE:** This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.

Date Shipment Needed: \_\_\_\_\_ Ship To:  Patient  Prescriber

Nursing needed;  Training needed ► All the supplies including syringes and needles will be dispensed if needed.

**DERMATOLOGY REFERRAL FORM ( L - Sil )**

**PATIENT INFORMATION**

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Sex:  M  F  Other: \_\_\_\_\_ Weight: \_\_\_\_\_  lbs.  kg.

SSN: \_\_\_\_\_ Phone: \_\_\_\_\_ Allergies: \_\_\_\_\_

Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_

Emergency Contact: \_\_\_\_\_ Phone: \_\_\_\_\_  **Additional Information Attached**

**PRESCRIBER INFORMATION**

Prescriber: \_\_\_\_\_ NPI: \_\_\_\_\_ DEA: \_\_\_\_\_ State Lic: \_\_\_\_\_

Supervising Physician: \_\_\_\_\_ Practice Name: \_\_\_\_\_

Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_

Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Key Office Contact: \_\_\_\_\_ Phone: \_\_\_\_\_

**DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT**

**Primary Diagnosis:**  L20, L20.8, L20.9 Atopic Dermatitis  L28.1 Prurigo nodularis  L40.0 Psoriasis  L40.1; L40.2; L40.3, L40.4, L40.8, L40.54 Psoriatic arthritis  L40.59  L50.1 Chronic Idiopathic Urticaria  L73.2 Hidradenitis Suppurativa  Other: \_\_\_\_\_

• Location:  Hands  Feet  Face  Scalp  Groin  Nails  Other: \_\_\_\_\_

• Severity:  Mild (up to 3% BSA)  Moderate (3-10% BSA)  Severe (greater than 10% BSA), BSA \_\_\_\_\_%

If treated previously for this condition, please indicate which drugs have been tried and failed: \_\_\_\_\_

Date range of previous therapy: \_\_\_\_\_

• Is patient currently on therapy?  Yes  No Type / medication(s): \_\_\_\_\_

• Will patient stop taking the above medication(s) before starting the new medication?  Yes  No, if yes, how long should patient wait before starting the new medication? \_\_\_\_\_

• Has patient received a PPD (tuberculosis) Skin Test?  Yes  No Results: \_\_\_\_\_

Prior to initiating treatment and periodically during therapy, patient should be evaluated for active tuberculosis and tested for latent infection.

**INSURANCE INFORMATION**

**Please attach front and back of patient's insurance card (medical and prescription)**

**COPY CARD ENROLLMENT**

**Please check if enrolling in copay card**      **Copay ID:** \_\_\_\_\_

**PRESCRIPTION INFORMATION**

**Nemluvio®**  **30 mg Pen** - The recommended subcutaneous dosage of NEMLUVIO for adult patients weighing less than 90 kg is an initial dose of 60 mg (two 30 mg injections), followed by 30 mg given every 4 weeks (Q4W)

Adult Patients Weighing less than 90 kg:  Inject 60 mg SQ once for initial dose      QTY: #2 Pens/28 DS | Refills: 0

Inject 30 mg SQ every 4 weeks      QTY: #1 Pens/28 DS | Refills: \_\_\_\_\_

**Nemluvio®**  **60 mg Pen** - The recommended subcutaneous dosage of NEMLUVIO for adult patients weighing 90 kg or more is an initial dose of 60 mg (two 30 mg injections), followed by 60 mg given every 4 weeks (Q4W)

Adult Patients Weighing 90 kg or more:  Inject 60 mg SQ once for initial dose      QTY: #2 Pens/28 DS | Refills: 0

Inject 60 mg SQ every 4 weeks      QTY: #2 Pens/28 DS | Refills: \_\_\_\_\_

**Odomzo®**  **200 mg Capsule PO Once Daily**      QTY: 30 CAPS | Refills: \_\_\_\_\_

**Otezla® Tablets** Plaque Psoriasis - moderate to severe; **Note:** Initial dose titration is intended to reduce GI symptoms

— **Adults:**

**Titration** DAY 1: 10 mg in morning; DAY 2: 10 mg in morning & 10 mg in evening; DAY 3: 10 mg in morning & 20 mg in evening;      QTY: 1 Month | Refills: 0

Dose: DAY 4: 20 mg in morning & 20 mg in evening; DAY 5: 20 mg in morning & 30 mg in evening; DAY 6 & thereafter: 30 mg twice daily

**Maintenance Dose:** 30 mg twice daily      QTY: 60 TABS (30 mg) | Refills: \_\_\_\_\_

**Other:** \_\_\_\_\_      QTY: \_\_\_\_\_ | Refills: \_\_\_\_\_

— **Children & Adolescents:** ≥ 6 years weighing 20 to < 50 kg:

**Otezla® 28 Day Treatment Initiation Pack:** DAY 1: Oral: 10 mg once daily in the morning; DAY 2: Oral: 10 mg twice daily;      QTY: #1 | Refills: 0

DAY 3: Oral: 10 mg in the morning & 20 mg in the evening; DAY 4 & thereafter: Oral: 20 mg twice daily

**Otezla® 20 mg Tablet:** Take 20 mg PO twice daily      QTY: 60 TABS | Refills: \_\_\_\_\_

— **Children & Adolescents:** ≥ 6 years weighing ≥ 50 kg:

**Otezla® 28 Day Treatment Initiation Pack:** DAY 1: Oral: 10 mg once daily in the morning; DAY 2: Oral: 10 mg twice daily;      QTY: #1 | Refills: 0

DAY 3: Oral: 10 mg in the morning & 20 mg in the evening; DAY 4: Oral: 20 mg twice daily;

DAY 5: Oral: 20 mg in the morning & 30 mg in the evening; DAY 6 & thereafter: Oral: 30 mg twice daily

**Otezla® 30 mg Tablet:** Take 30 mg PO twice daily      QTY: 60 TABS | Refills: \_\_\_\_\_

**Remicade®** 100 mg Vial     **Inflectra®** 100 mg Powder Vial     **Renflexis®** 100 mg Powder Vial     **Avsola®** 100 mg Powder Vial       **Enroll in AccessOneSM Program**

MD's Office Infusion     Home Infusion Supplies Required       Starter Dose not needed

**Starter Dose:** \_\_\_\_\_ mg IV on Week 0, Week 2, Week 6, then      QTY: QS 3 Infusions | Refills: 0

**Maintenance Dose:** \_\_\_\_\_ mg IV every \_\_\_\_\_ weeks      QTY: QS 1 Infusion | Refills: \_\_\_\_\_

**Rinvoq®**  **15 mg Tablet**     **30 mg Tablet**    Take 1 tablet PO once daily      QTY: 1 Month | Refills: \_\_\_\_\_

**Siliq®**  **210 mg / 1.5 mL Prefilled Syringe (2 pack)**       Starter Dose not needed       **Enroll in REMS Program**

**Starter Dose for Plaque Psoriasis:** 210 mg SQ at Weeks 0, 1 and 2, followed by maintenance dose      QTY: 1 Box (2 PFS) | Refills: 0

**Maintenance Dose for Plaque Psoriasis:** 210 mg SQ once every two weeks (Starting at Week 2)      QTY: 1 Box (2 PFS) | Refills: \_\_\_\_\_

**Prescriber's Signature:** \_\_\_\_\_  DAW (Dispense as Written)      **Date:** \_\_\_\_\_

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. NO STAMPED SIGNATURES WILL BE ACCEPTED. Where required by law, send electronic prescription or on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

**IMPORTANT NOTICE:** This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.

Date Shipment Needed: \_\_\_\_\_ Ship To:  Patient  Prescriber

Nursing needed;  Training needed ► All the supplies including syringes and needles will be dispensed if needed.

**DERMATOLOGY REFERRAL FORM ( Sim - Z )**

**PATIENT INFORMATION**

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Sex:  M  F  Other: \_\_\_\_\_ Weight: \_\_\_\_\_  lbs.  kg.

SSN: \_\_\_\_\_ Phone: \_\_\_\_\_ Allergies: \_\_\_\_\_

Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_

Emergency Contact: \_\_\_\_\_ Phone: \_\_\_\_\_  Additional Information Attached

**PRESCRIBER INFORMATION**

Prescriber: \_\_\_\_\_ NPI: \_\_\_\_\_ DEA: \_\_\_\_\_ State Lic: \_\_\_\_\_

Supervising Physician: \_\_\_\_\_ Practice Name: \_\_\_\_\_

Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_

Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Key Office Contact: \_\_\_\_\_ Phone: \_\_\_\_\_

**DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT**

**Primary Diagnosis:**  L20, L20.8, L20.9 Atopic Dermatitis  L28.1 Prurigo nodularis  L40.0 Psoriasis  L40.1; L40.2; L40.3, L40.4, L40.8, L40.54 Psoriatic arthritis  L40.59  L50.1 Chronic Idiopathic Urticaria  L73.2 Hidradenitis Suppurativa  Other: \_\_\_\_\_

• Location:  Hands  Feet  Face  Scalp  Groin  Nails  Other: \_\_\_\_\_

• Severity:  Mild (up to 3% BSA)  Moderate (3-10% BSA)  Severe (greater than 10% BSA), BSA \_\_\_\_\_ %

If treated previously for this condition, please indicate which drugs have been tried and failed: \_\_\_\_\_

Date range of previous therapy: \_\_\_\_\_

• Is patient currently on therapy?  Yes  No Type / medication(s): \_\_\_\_\_

• Will patient stop taking the above medication(s) before starting the new medication?  Yes  No, if yes, how long should patient wait before starting the new medication? \_\_\_\_\_

• Has patient received a PPD (tuberculosis) Skin Test?  Yes  No Results: \_\_\_\_\_

Prior to initiating treatment and periodically during therapy, patient should be evaluated for active tuberculosis and tested for latent infection.

**INSURANCE INFORMATION**

Please attach front and back of patient's insurance card (medical and prescription)

**COPAY CARD ENROLLMENT**

Please check if enrolling in copay card **Copay ID:** \_\_\_\_\_

**PRESCRIPTION INFORMATION**

**Simponi®**  Aria 50 mg / 4 mL Patient Weight (kg): \_\_\_\_\_  Starter Dose not needed  Enroll in SimponiOne® Program

Starter Dose: 2 mg / kg IV at Weeks 0 and 4 QTY: \_\_\_\_\_ 1 Month | Refills: 0

Maintenance Dose: 2 mg / kg IV every 8 weeks QTY: \_\_\_\_\_ QS for 8 Weeks | Refills: \_\_\_\_\_

**Simponi®**  SmartJect 50 mg / 0.5 mL  50 mg / 0.5 mL Prefilled Syringe \*Pens will be dispensed if no preference is indicated

50 mg SQ every month QTY: \_\_\_\_\_ 1 Month | Refills: \_\_\_\_\_

Other: \_\_\_\_\_ QTY: \_\_\_\_\_ | Refills: \_\_\_\_\_

**Skyrizi®**  150 mg / mL Pen Autoinjector  150 mg / mL Prefilled Syringe \*Pens will be dispensed if no preference is indicated

Starter Dose: 150 mg SQ at Week 0 and 4  Starter Dose not needed QTY: \_\_\_\_\_ 1 | Refills: 1

Maintenance Dose: 150 mg SQ every 12 weeks QTY: \_\_\_\_\_ 1 | Refills: \_\_\_\_\_

**Sotyku®**  6 mg PO once daily QTY: \_\_\_\_\_ | Refills: \_\_\_\_\_

**Stelara®**  Prefilled Syringe  Vial  MD's Office Infusion  Home Infusion Supplies Required \*Prefilled Syringe will be dispensed if preference is not indicated

≤ 100 kg Starter Dose: 45 mg SQ initially (Week 0), then 45 mg SQ after 4 weeks of initial dose (Week 4)  Enroll in Janssen CarePath Program

≤ 100 kg Maintenance Dose: 45 mg SQ every 12 weeks  Other: \_\_\_\_\_ QTY: \_\_\_\_\_ 1 x 45 mg | Refills: 1

> 100 kg Starter Dose: 90 mg SQ initially (Week 0), then 90 mg SQ after 4 weeks of initial dose (Week 4) QTY: \_\_\_\_\_ 1 x 45 mg | Refills: \_\_\_\_\_

> 100 kg Maintenance Dose: 90 mg SQ every 12 weeks  Other: \_\_\_\_\_ QTY: \_\_\_\_\_ 1 x 90 mg | Refills: 1

> 100 kg Maintenance Dose: 90 mg SQ every 12 weeks  Other: \_\_\_\_\_ QTY: \_\_\_\_\_ 1 x 90 mg | Refills: \_\_\_\_\_

**Taltz®**  80 mg / mL Autoinjector  80 mg / mL Prefilled Syringe \*Pen will be dispensed if no preference is indicated  Starter Dose not needed

Starter Dose for Plaque Psoriasis: 160 mg (two 80 mg inj.) at Week 0, then 80 mg at Week 2, 4, 6, 8, 10, 12 QTY: \_\_\_\_\_ 8 | Refills: 0

Maintenance Dose for Plaque Psoriasis: 80 mg every 4 weeks QTY: \_\_\_\_\_ 1 | Refills: \_\_\_\_\_

Starter Dose for Psoriatic Arthritis: 160 mg (two 80 mg inj.) at Week 0 QTY: \_\_\_\_\_ 2 | Refills: 0

Maintenance Dose for Psoriatic Arthritis: 80 mg every 4 weeks QTY: \_\_\_\_\_ 1 | Refills: \_\_\_\_\_

Other: \_\_\_\_\_ QTY: \_\_\_\_\_ | Refills: \_\_\_\_\_

**Tremfya®**  100 mg / mL Pen Autoinjector  100 mg / mL Prefilled Syringe \*Pens will be dispensed if no preference is indicated

Starter Dose: 100 mg SQ at Week 0, 4, and every 8 weeks thereafter  Starter Dose not needed QTY: \_\_\_\_\_ 1 | Refills: 0

Maintenance Dose: 100 mg SQ every 8 weeks (starting at Week 4) QTY: \_\_\_\_\_ 1 | Refills: \_\_\_\_\_

**Keljanz®**  5 mg Tablet  10 mg Tablet: 1 tablet PO twice daily QTY: \_\_\_\_\_ 1 Month | Refills: \_\_\_\_\_

11 mg ER Tablet: 1 tablet PO daily QTY: \_\_\_\_\_ 1 Month | Refills: \_\_\_\_\_

**Xolair®**  150 mg Prefilled Syringe  150 mg Vial

150 mg SQ every 4 weeks  300 mg SQ every 4 weeks QTY: \_\_\_\_\_ 28 Day Supply | Refills: \_\_\_\_\_

**Prescriber's Signature:** \_\_\_\_\_  DAW (Dispense as Written) **Date:** \_\_\_\_\_

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. NO STAMPED SIGNATURES WILL BE ACCEPTED. Where required by law, send electronic prescription or on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

**IMPORTANT NOTICE:** This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.