

Date Shipment Needed: _____ Ship To: Patient Prescriber

HEMOPHILIA AND CLOTTING DISORDER REFERRAL FORM

PATIENT INFORMATION					
Patient Name:		DOB:	Sex: <input type="checkbox"/> M <input type="checkbox"/> F <input type="checkbox"/> Other:	Weight:	<input type="checkbox"/> lbs. <input type="checkbox"/> kg.
SSN:	Phone:	Allergies:			
Address:			City:	State:	Zip:
Emergency Contact:		Phone:	<input type="checkbox"/> Additional Information Attached		

PRESCRIBER INFORMATION					
Prescriber:		NPI:	DEA:	State Lic:	
Supervising Physician:			Practice Name:		
Address:		City:	State:	Zip:	
Phone:	Fax:	Key Office Contact:		Phone:	

DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT	
Primary Diagnosis: (ICD-10 Code & Description) <input type="checkbox"/> D66 Hereditary factor VIII deficiency <input type="checkbox"/> D67 Hereditary factor IX deficiency <input type="checkbox"/> D68.0 Von Willebrand's disease <input type="checkbox"/> D68.311 Acquired Hemophilia <input type="checkbox"/> D68.8 Other specified coagulation defects <input type="checkbox"/> D68.9 Coagulation defect, unspecified <input type="checkbox"/> D68.2 Hereditary deficiency of other clotting factors Other Code: _____ Description: _____ Other medications patient is currently taking including OTC medications with dosage and direction (or fax medication profile): _____	

INSURANCE INFORMATION	
<input type="checkbox"/> Please attach front and back of patient's insurance card (medical and prescription)	

COPAY CARD ENROLLMENT	
<input type="checkbox"/> Please check if enrolling in copay card	Copay ID: _____

NURSING	
<input type="checkbox"/> Skilled Nursing Needed: <input type="checkbox"/> No <input type="checkbox"/> Yes (NOTE: not available for PRN or pre-op dosing) As needed for IV access, administration, and proper clinical monitoring. If yes, specify services needed: <input type="checkbox"/> Teach & Train (up to 3 visits) <input type="checkbox"/> Post-Op (Must include date of procedure: _____) Site of Care: <input type="checkbox"/> MD office <input type="checkbox"/> Infusion Clinic <input type="checkbox"/> Outpatient Health <input type="checkbox"/> Home Health	

PRESCRIPTION INFORMATION				
FACTOR REPLACEMENT	PROPHYLAXIS	BREAKTHROUGH BLEEDS	BREAKTHROUGH BLEEDS	REFILLS
<input type="checkbox"/> Advate <input type="checkbox"/> Eloclate <input type="checkbox"/> Nuwiq <input type="checkbox"/> Adynovate <input type="checkbox"/> Hemofil-M <input type="checkbox"/> Recombinate <input type="checkbox"/> Afstyla <input type="checkbox"/> Humate-P* <input type="checkbox"/> Rixubis <input type="checkbox"/> Alphanate* <input type="checkbox"/> Idelvion <input type="checkbox"/> Vonvendi <input type="checkbox"/> Alphanine <input type="checkbox"/> Ixinity <input type="checkbox"/> Wilate* <input type="checkbox"/> Alprolix <input type="checkbox"/> Jivi <input type="checkbox"/> Xyntha <input type="checkbox"/> Altuviiio <input type="checkbox"/> Kovaltry <input type="checkbox"/> Other: _____ <input type="checkbox"/> Benefix <input type="checkbox"/> Novoeight	Dose: _____ units Assay Variance: +/- 10% or <input type="checkbox"/> Other: _____ Route: IV Frequency: _____ Dose Quantity: _____ <small>* For these products, indicate if doses are expressed as Factor VIII or VWF:RCoF units</small>	Bleed Type: <input type="checkbox"/> Major <input type="checkbox"/> Minor <input type="checkbox"/> Other: _____ Dose: _____ units Assay Variance: +/- 10% or <input type="checkbox"/> Other: _____ Route: IV Frequency: _____ prn Dose Quantity: _____	Bleed Type: <input type="checkbox"/> Major <input type="checkbox"/> Minor <input type="checkbox"/> Other: _____ Dose: _____ units Assay Variance: +/- 10% or <input type="checkbox"/> Other: _____ Route: IV Frequency: _____ prn Dose Quantity: _____	11 or _____

Alhemo (Hemophilia A&B with inhibitors) — Loading dose: Inject 1 mg/kg (Day 1) SQ x 1; Maintenance dose: Inject 0.2 mg/kg SQ daily x 4 weeks; (measure Concizimab-mtci plasma conc)
 Individualized Maintenance Dose: Inject _____ mg/kg SQ daily; Refills: _____;
 Hympavzi (Hemophilia A&B without inhibitors) — Loading dose: Inject 300 mg SQ x 1; Maintenance dose: Inject 150 mg SQ weekly; Refills: _____

HEMLIBRA	LOADING DOSE	MAINTENANCE DOSE	BREAKTHROUGH BLEEDS	REFILLS
<input type="checkbox"/> Hemlibra 30 mg/mL (will require separate injection) <input type="checkbox"/> Hemlibra 60 mg/0.4 mL <input type="checkbox"/> Hemlibra 105 mg/0.7 mL <input type="checkbox"/> Hemlibra 150 mg/mL May use any combination of available strengths to dispense calculated dose, unless otherwise requested	<input type="checkbox"/> No Loading Dose Needed <input type="checkbox"/> 3 mg/kg SQ once weekly for 4 weeks Calculated Dose: _____ mg Dose Quantity: 4 ; Refills: 0	<input type="checkbox"/> 1.5 mg/kg SQ every week <input type="checkbox"/> 3.0 mg/kg SQ every 2 weeks <input type="checkbox"/> 6.0 mg/kg SQ every 4 weeks Calculated Dose: _____ mg Dose Quantity: _____ ; Refills: _____	Please select a Factor VIII product in the section above for treatment of breakthrough bleeds while on Hemlibra	11 or _____

ANCILLARY MEDICATIONS / SUPPLIES			
PRODUCT	DIRECTIONS	QUANTITY	REFILLS
<input type="checkbox"/> Sodium chloride 0.9% flush syringe (10 mL)	Access Device: <input type="checkbox"/> Port <input type="checkbox"/> PICC <input type="checkbox"/> PIV <input type="checkbox"/> Butterfly Flush with _____ mL every _____	<input type="checkbox"/> 1 month <input type="checkbox"/> 3 months <input type="checkbox"/> Other: _____	11 or _____
<input type="checkbox"/> Heparin 10 units/mL flush syringe (5 mL) <input type="checkbox"/> Heparin 100 units/mL flush syringe (5 mL)	Access Device: <input type="checkbox"/> Port <input type="checkbox"/> PICC <input type="checkbox"/> PIV <input type="checkbox"/> Butterfly Flush with _____ mL every _____	<input type="checkbox"/> 1 month <input type="checkbox"/> 3 months <input type="checkbox"/> Other: _____	11 or _____
<input type="checkbox"/> Needles and Syringes	Use as directed for administration of infused/injected medication	<input type="checkbox"/> 1 month <input type="checkbox"/> 3 months <input type="checkbox"/> Other: _____	11 or _____
Epinephrine is Required for Nursing <input type="checkbox"/> Epinephrine 0.3 mg auto-injector (Adult: > 30 kg) <input type="checkbox"/> Epinephrine 0.15 mg auto-injector (15-30 kg)	Inject 1 pen IM PRN severe allergic reaction – Call 911 May repeat in 5-15 minutes as needed	<input type="checkbox"/> 1 Box (2 Pens) <input type="checkbox"/> Other: _____	11 or _____
<input type="checkbox"/> Lidocaine/Prilocaine 2.5% / 2.5% cream	Apply to injection site as needed Day Supply: _____	<input type="checkbox"/> 1 Tube (30 grams) <input type="checkbox"/> Other: _____	11 or _____
<input type="checkbox"/> Other Product: _____	Directions: _____ (Include dose, route and frequency)	Quantity: _____	11 or _____

Prescriber's Signature: _____ **Date:** _____

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. NO STAMPED SIGNATURES WILL BE ACCEPTED. Where required by law, send electronic prescription or on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

IMPORTANT NOTICE: This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.